arXiv:0707.3285v3 [hep-ph] 21 Dec 2007

WSU-HEP-0704

On the Non-Manifest Left-Right Model Contribution to the Lifetime Difference in $D^0 - \bar{D}^0$ Mixing.

G. K. Yeghiyan[*]

Department of Physics and Astronomy
Wayne State University, Detroit MI, USA.

Abstract

New physics contribution to the lifetime difference in $D^0 - \bar{D}^0$ mixing is re-examined within the non-manifest Left-Right Symmetric Model. Diagrams with one of $\Delta C = 1$ transitions, mediated by a propagator with $W_L - W_R$ mixing, are revisited. While these diagrams are believed to give the dominant contribution, compatible with the experimental data, it is shown that due to GIM cancelation, such diagrams are negligible in sum. Thus, Left-Right Symmetric Model contribution to the lifetime difference in $D^0 - \bar{D}^0$ mixing is about two orders of magnitude less than actual experimental value for $\Delta\Gamma_D$.

Observation of $D^0 - \bar{D}^0$ oscillations by BaBaR and Belle collaborations [1, 2] has revived the theoretical interest to this phenomenon. In particular, new physics contribution to $D^0 - \bar{D}^0$ mixing has been re-considered in several publications [3] - [13].

The impact of a new physics on the lifetime difference in $D^0 - \bar{D}^0$ mixing has been in details studied in [10]. Model independent analysis has been performed and the derived analytical formulae have been then applied within several extensions of the Standard Model. It has been shown that to the lowest order in the perturbation theory, new physics contribution to $\Delta\Gamma_D$ may be several orders of magnitude greater than that of the Standard Model. Later on the lifetime difference in $D^0 - \bar{D}^0$ mixing has been also considered in [11, 12] and [13], within the supersymmetric models with R-parity violation and the Left-Right Symmetric Model, respectively.

It has been argued in [13] that within the non-manifest Left-Right (LR) Symmetric Model, new physics contribution to the lifetime difference in $D^0 - \bar{D}^0$ mixing may be significant:

$$|y_{LR}| \equiv \frac{|\Delta\Gamma_{D_{LR}}|}{2\Gamma_D} \leq 1.4 \times 10^{-3}, \qquad (1)$$

which means that y_{LR} may be of the same order as the experimental value of y [14],

$$y \equiv \frac{\Delta\Gamma_D}{2\Gamma_D} = (6.6 \pm 2.1) \times 10^{-3} \qquad (2)$$

This result has been derived by considering the box diagrams with one of $\Delta C = 1$ transitions being generated by a new physics (NP) interaction and mediated by a propagator with $W_L - W_R$ mixing (Fig. 1). Note that W_R part of the propagator couples with the u-quark, which allows one to remove a power of the suppression in terms of $\lambda = \sin\theta_C \approx 0.23$.



FIG. 1: $\Delta C = 1$ transition mediated by a propagator with $W_L - W_R$ mixing.

In this letter we revisit the contribution of the box diagrams with the new physics generated $\Delta C = 1$ transition, presented in Fig 1. While the analysis of ref. [13] is restricted by considering only the diagrams with the intermediate s-quark states, i.e. $q = s$ and $q' = s$, we include also the diagrams with $q = d$ and/or $q' = d$. We will see that diagrams with the intermediate d-quark states may not be neglected, in spite of $m_d \ll m_s$. Moreover, they play crucial role in taking into account GIM cancelation effects properly.

In this paper we show that box diagrams with the new physics generated $\Delta C = 1$ transition, presented in Fig 1, are negligible *in sum* due to GIM cancelation. Thus, one must replace the bound on y_{LR}, given by equation (1), by

---

[*] Electronic address: ye˙gagik@wayne.edu

$$|y_{LR}| \leq 8.8 \times 10^{-5} \tag{3}$$

This constraint on y_{LR} has been derived in [10], neglecting $\Delta C = 1$ transition presented in Fig. 1.

For $\Delta C = 1$ interaction in Fig. 1, the relevant part of the low-energy effective Hamiltonian has the following form:

$$H^{\Delta C=1}_{W_{L-R}} = -\frac{4G_F\xi_g}{\sqrt{2}} \sum_{q,q'} V^{*L}_{cq} V^{R}_{uq'} \Big[\bar{C}_1(m_c)Q_1 + +\bar{C}_2(m_c)Q_2\Big] \tag{4}$$

$$Q_1 = \bar{u}_i\gamma^\nu P_R q'_j \bar{q}_j\gamma_\nu P_L c_i, \quad Q_2 = \bar{u}_i\gamma^\nu P_R q'_i \bar{q}_j\gamma_\nu P_L c_j$$

where $P_L = (1-\gamma_5)/2$, $P_R = (1+\gamma_5)/2$, i, j stand for color indices, V^L, V^R are the left- and right-handed quark CKM matrices and ξ_g is defined in [13]. If only one $\Delta C = 1$ transition in the box diagrams is generated by an NP interaction, the approach described in ref. [10] may be used. For the new physics $\Delta C = 1$ effective Hamiltonian given by equation (4), it is not hard to see that only the term $I_4(x_q, x_{q'})\ \langle\ \bar{D}^0 \mid O_4^{ijkl} \mid D^0\ \rangle$ in equation (7) of [10] contributes. Basically, this result is in agreement with that of ref. [13], however there is an essential difference. While $q = q' = s$ in [13], we take here $q = s, d$ and $q' = s, d$. If one denotes by $y^{(1)}_{LR}$ the considered here contribution to the lifetime difference in $D^0 - \bar{D}^0$ mixing, then, using eqs. (7), (9), (10) in ref. [10] (setting there $D_{qq'} = -\left(G_F/\sqrt{2}\right)\xi_g V^{L*}_{cq} V^R_{uq'}$, $\bar{\Gamma}_1 = \gamma^\nu P_R$, $\bar{\Gamma}_2 = \gamma_\nu P_L$), it is straightforward to show after doing some algebra that

$$y^{(1)}_{LR} = \sum_{q,\ q'} C^{qq'}_{LR}\ V^{L^*}_{cq'} V^R_{uq'}\ \Big[K_2\langle Q'\rangle + K_1\langle\tilde{Q}'\rangle\Big] \tag{5}$$

where

$$C^{qq'}_{LR} = \frac{G_F^2 m_c^2\ \xi_g}{2\pi m_D \Gamma_D}\ V^{L^*}_{cq} V^L_{uq}\ \sqrt{x_{q'}}\ \Big[(1-x_{q'})^2 - -2x_q x_{q'} - x_q^2\Big] \tag{6}$$

and the notations in (5) and (6) are the same as in [13].

Formulae (5) and (6) are generalization of formulae (3) and (4) of ref. [13] for the case when both s- and d-quark intermediate states are considered, thus C_{LR} of [13] is replaced here by $C^{qq'}_{LR}$ and sum over q, q' is implemented. Else, in order to take properly into account GIM cancelation effects, we keep in equation (6) higher order terms in the expansion in powers of $x_q \equiv m_q^2/m_c^2$ and $x_{q'} \equiv m_{q'}^2/m_c^2$.

It is worth to note that dependence on x_q appears only in the next-to-next-to-leading order terms of this expansion. The difference in the behavior of $y^{(1)}_{LR}$ with x_q and with $x_{q'}$ is related to different chiralities of the light quarks q and q' in (4). More detailed discussion of the behavior of $D^0 - \bar{D}^0$ mixing amplitude with the light quark masses, depending on these quarks chiralities, may be found in refs. [15]-[17]. Discussion for a particular case of the width difference is also available in [18, 19].

It is clear from (6) that if one takes the limit $x_d \equiv m_d^2/m_c^2 = 0$, $C^{qq'}_{LR} = 0$ for $q' = d$. Thus, formula (5) is significantly simplified:

$$y^{(1)}_{LR} = \left[C^{ss}_{LR} + C^{ds}_{LR}\right]\ V^{L^*}_{cs} V^R_{us}\ \Big[K_2\langle Q'\rangle + K_1\langle\tilde{Q}'\rangle\Big] \tag{7}$$

where

$$C^{ss}_{LR} = \frac{G_F^2 m_c^2\ \xi_g}{2\pi m_D \Gamma_D}\ V^{L^*}_{cs} V^L_{us}\ \sqrt{x_s}\ \Big[(1-x_s)^2 - 3x_s^2\Big] \tag{8}$$

$$C^{ds}_{LR} = \frac{G_F^2 m_c^2\ \xi_g}{2\pi m_D \Gamma_D}\ V^{L^*}_{cd} V^L_{ud}\ \sqrt{x_s}\ (1-x_s)^2 \tag{9}$$

As it follows from (7) - (9), in the limit $m_d = 0$ there is an additional contribution - as compared to that of ref. [13] - from the diagram in Fig. 1 when $q = d$ and $q' = s$: $C^{ds}_{LR} \neq 0$. Moreover, using the fact that $V^{L^*}_{cs} V^L_{us} \approx -V^{L^*}_{cd} V^L_{ud} + O(\lambda^5)$, it is not hard to see that $C^{ss}_{LR} \approx -C^{ds}_{LR}$ with accuracy of the terms $\sim \lambda^5$ or $\sim x_s^{5/2}$. Thus, sum of C^{ss}_{LR} and C^{ds}_{LR} is much less in the absolute value than these quantities by themselves. This is manifestation of (approximate) GIM cancelation that makes $y^{(1)}_{LR}$ negligible.

Using the unitarity condition,

$$V^{L^*}_{cs} V^L_{us} + V^{L^*}_{cd} V^L_{ud} + V^{L^*}_{cb} V^L_{ub} = 0 \tag{10}$$

one gets after doing some algebra

$$C^{ss}_{LR} + C^{ds}_{LR} = \frac{G_F^2 m_c^2\ \xi_g}{2\pi m_D \Gamma_D}\ \sqrt{x_s}\ \Big[-Re\left(V^{L^*}_{cb} V^L_{ub}\right)(1 - -x_s)^2 -\ 3\ V^{L^*}_{cs} V^L_{us}\ x_s^2\Big] \tag{11}$$

Note that unlike CKM products in (5) - (9), $V^{L^*}_{cb} V^L_{ub}$ has non-negligible phase [20], thus one must explicitly indicate that the real part of this product is only relevant. It is assumed no new source of CP-violation [10] (V_R is real and no spontaneous CP-violation). In this case, the impact of CP-violating effects on $\Delta\Gamma_D$ is negligible.

Usually, when studying $D^0 - \bar{D}^0$ oscillations, one puts $V^{L^*}_{cb} V^L_{ub} \approx 0$, as $|V^{L^*}_{cb} V^L_{ub}| \ll V^{L^*}_{cs} V^L_{us}$. This is the two quark generation mixing approximation, that is widely applied in studying $D^0 - \bar{D}^0$ mixing effects within the Standard Model (see e.g [21]) and some of its extensions. However, in our case this approximation is not valid. Indeed, using $Re\left(V_{cb}^{L^*} V_{ub}^L\right) \approx A^2 \lambda^5 \rho$ and $V_{cs}^{L^*} V_{us}^L \approx \lambda$, it is not hard to see that the first term in the square brackets in (11) dominates over the last one, for $A \approx 0.82$, $\lambda \approx 0.23$, $\rho \approx 0.23$ [20] and $x_s \equiv m_s^2(m_c)/m_c^2(m_c) \approx 0.007$ [12].

To the lowest order in the perturbation theory, one gets a rough estimate of the effect rather than a precise numerical evaluation. In what follows, one may to a good approximation disregard the subdominant terms in (11). Then, one may rewrite equation (7) in a more compact form:

$$y_{LR}^{(1)} = -\bar{C}_{LR}\, V_{cs}^{L^*} V_{us}^R \left[K_2 \langle Q' \rangle + K_1 \langle \tilde{Q}' \rangle\right] \qquad (12)$$

where

$$\bar{C}_{LR} = \frac{G_F^2 m_c^2\, \xi_g}{2\pi m_D \Gamma_D}\, Re\left(V_{cb}^{L^*} V_{ub}^L\right) \sqrt{x_s} \qquad (13)$$

We parameterize $\langle Q' \rangle$ and $\langle \tilde{Q}' \rangle$, using the moderate vacuum saturation approach [4]:

$$\begin{aligned} \langle Q' \rangle &\equiv \langle \bar{D}^0 | \bar{u}_i \gamma^\mu P_L c_i \bar{u}_j \gamma_\mu P_R c_j | D^0 \rangle = \\ &= -\frac{1}{2} f_D^2 m_D^2 B_D - \frac{1}{3} f_D^2 m_D^2 \left(\frac{m_D}{m_c}\right)^2 B_D^S \qquad (14) \\ \langle \tilde{Q}' \rangle &\equiv \langle \bar{D}^0 | \bar{u}_i \gamma^\mu P_L c_j \bar{u}_j \gamma_\mu P_R c_i | D^0 \rangle = \\ &= -\frac{1}{6} f_D^2 m_D^2 B_D - f_D^2 m_D^2 \left(\frac{m_D}{m_c}\right)^2 B_D^S \qquad (15) \end{aligned}$$

where $f_D \approx 0.22$GeV [22], $B_D \approx 0.8$ [23], and we choose $B_D^S \approx B_D$. Then, using $G_F = 1.166 \times 10^{-5} GeV^{-2}$, $\Gamma_D \approx 1.6 \times 10^{-12}$GeV, $m_D \approx 1.865$GeV, $m_c \equiv m_c(m_c) \approx 1.25$GeV [20], $K_1 \equiv 3C_1\tilde{C}_1 + C_1\tilde{C}_2 + C_2\tilde{C}_1 \approx 3C_1^2 + 2C_1C_2$, $K_2 \equiv C_2\tilde{C}_2 \approx C_2^2$, $C_1(m_c) = -0.411$, $C_2(m_c) = 1.208$ [21], $V_{cs}^L \approx 1 - \lambda^2/2$, $V_{us}^R \approx 1$ and [13, 24] $\xi_g \leq 0.033$, one gets

$$y_{LR}^{(1)} \leq 2.6 \times 10^{-7} \qquad (16)$$

Thus, due to GIM cancelation, box diagrams with the new physics generated $\Delta C = 1$ transition, presented in Fig. 1, give *in sum* negligible contribution to the lifetime difference in $D^0 - \bar{D}^0$ mixing.

It is left for a reader to verify that one gets negligible contribution to y_{LR} also in the case when $W_L - W_R$ propagator in Fig. 1 is flipped so that W_R couples with the charm quark.

In what follows, one should use the result of ref. [10] that has been derived neglecting $\Delta C = 1$ transition in Fig. 1. In other words, one should use the bound on y_{LR}, given by equation (3). Thus, within the non-manifest Left-Right Symmetric Model, new physics contribution to to the lifetime difference in $D^0 - \bar{D}^0$ mixing is rather small.

In is worth to note here that this result has been derived considering the diagrams with only one $\Delta C = 1$ transition generated by a new physics interaction. There are also box diagrams with both $\Delta C = 1$ transitions occurring due to NP interactions. These diagrams have not been considered so far, as within the Left-Right Symmetric Model they are estimated to have a small contribution to $\Delta\Gamma_D$. On the other hand, it is still possible that within the non-manifest version of the LR model, there are some corners of the parameter space with M_{W_R} below 1 TeV [25], where such diagrams are perhaps non-negligible. Study of this possibility requires detailed and careful scanning of the parameter space of the theory, taking into account all possible constraints, coming from $K_L - K_S$ and $B^0 - \bar{B}^0$ mass differences, as well as other phenomenological constraints. Such a detailed analysis is out of the scope of this brief report.

In conclusion, lifetime difference in $D^0 - \bar{D}^0$ oscillations has been re-considered within the non-manifest Left-Right Symmetric Model. It has been shown that, due to GIM cancelation effects, new physics contribution to the lifetime difference in $D^0 - \bar{D}^0$ mixing is rather small, as compared to the experimental value of $\Delta\Gamma_D$.

Author is grateful to A. A. Petrov and Chuan-Hung Chen for valuable discussions. This work has been supported by the grants NSF PHY-0547794 and DOE DE-FGO2-96ER41005.

---

[1] The BaBaR Colaboration, B. Aubert, et al. hep-ex/0705.0704;
The BaBaR Colaboration, B. Aubert, et al., hep-ex/0703020.

[2] Belle Collaboration, L. M. Zhang et al., hep-ex/0704.1000;
Belle Colaboration, M. Staric, et al., Phys. Rev. Lett. **98** (2007) 211803.

[3] M. Ciuchini et al., hep-ph/0703204.

[4] E. Golowich et al.,hep-ph/0705.3650.

[5] M. Blanke et al., hep-ph/0703254.
[6] X. G. He, G. Valencia, hep-ph/0703270.
[7] Ch. H. Chen, Ch. Q. Geng, T. Ch. Yuan, hep-ph/0704.0601.
[8] X. Q. Li, Z. T. Wei, hep-ph/0705.1821.
[9] B. Dutta, Y. Mimura, hep-ph/0708.3080.
[10] E. Golowich, S. Pakvasa, A. A. Petrov, Phys. Rev. Lett. **98** (2007) 181801.
[11] Sh. L. Chen et al., hep-ph/0706.1100.
[12] A. A. Petrov, G. K. Yeghiyan, hep-ph/0710.4939.
[13] Chuan-Hung Chen, Chao-Qiang Geng, Soo-Hyeon Nam, Phys. Rev. Lett. **99** (2007) 019101.
[14] Heavy Flavor Averaging Group, http://www.slac.stanford.edu/xorg/hfag/charm/index.html
[15] H. Georgi, Phys. Lett. **B 297** (1992) 353.
[16] T Ohl et al., Nucl. Phys. **B 403** (1993) 605.
[17] I. I. Bigi, N. G. Uraltsev, Nucl. Phys. **B 592** (2001) 92.
[18] A. F. Falk, Y. Grossman, Z. Ligeti and A. A. Petrov, Phys. Rev. **D 65** (2002) 054034.
[19] A. A. Petrov, Proceedings of 4th Workshop on Continuous Advances in QCD, Minneapolis, Minnesota, 12-14 May 2000, hep-ph/0009160.
[20] W. M. Yao et al. (Particle Data Group), Journal of Physics **G 33** (2006) 1.
[21] E. Golowich, A. A. Petrov, Phys.Lett. **B 625** (2005) 53.
[22] D. Asner, Contribution to WG2 Report on Flavor in the ERA of the LHC, CERN, March 26-28, 2007.
[23] R. Gupta, T. Bhattacharya, S. R. Sharpe, Phys. Rev. **D 55** (1997) 4036.
[24] S. H. Nam, Phys. Rev. **D 68** (2003)115006.
[25] P. Langacker, S. U. Sankar, Phys. Rev. **D 40** (1989) 1569.